

Bloch Electron on a Triangular Lattice and Quantum Ising Model in a Transverse Field

Jukka A. Ketoja

*Department of Physics, Åbo Akademi,
Porthansgatan 3, FIN-20500 Åbo, Finland*

Indubala I. Satija

*Department of Physics and
Institute of Computational Sciences and Informatics,
George Mason University,
Fairfax, VA 22030
(February 1, 2008)*

The tight binding model for an electron on an anisotropic triangular lattice in a uniform magnetic field is studied using a decimation scheme. The model exhibits a transition from critical to localized phase and the phase diagram is described in terms of three nontrivial renormalization fixed points for the band edges. These subcritical, critical, and supercritical universality classes also describe the corresponding states of the quantum Ising chain in a modulating transverse field. The only exception is the conformally invariant point of the Ising model which has no analog in the electron problem.

75.30.Kz, 64.60.Ak, 64.60.Fr

The problem of a two-dimensional Bloch electron in an irrational magnetic flux is important both theoretically and experimentally. [1] In this paper, we study the electron on a triangular lattice with anisotropic couplings between the nearest neighbor sites. We use a renormalization approach to obtain the phase diagram and universal scaling properties of the model. The results are compared with those of the Ising model in a transverse modulating field of periodicity incommensurate with the periodicity of the spin chain. With the exception of conformally invariant point [2] of the Ising model [3], the band edge as well as the band center states of the two models belong to the same universality class.

In the tight binding approximation, the energy spectrum for a single Bloch band of an anisotropic triangular lattice with couplings t_1 and t_2 is [4],

$$\epsilon(k_x, k_y) = -t_1 \cos\left[\frac{a}{2}(\sqrt{3}k_x - k_y)\right] - t_1 \cos\left[\frac{a}{2}(\sqrt{3}k_x + k_y)\right] + t_2 \cos(ak_y) \quad (1)$$

where a is the lattice spacing. Using Peierl's substitution $\hbar\mathbf{k} \rightarrow \mathbf{p} - \frac{e}{c}\mathbf{A}$ with $\mathbf{p} = \frac{\hbar}{i}\nabla$ and the Landau gauge $\mathbf{A} = (0, x, 0)H$ for the vector potential $\mathbf{A}$, the two-dimensional problem is reduced to the one-dimensional tight binding model (TBM) [4]

$$\cos[\pi(i\sigma + (\phi + \frac{\sigma}{2}))]\psi_{i+1} + \cos[\pi(i\sigma + (\phi - \frac{\sigma}{2}))]\psi_{i-1}$$

$$+ \lambda \cos[2\pi(i\sigma + \phi)]\psi_i = E\psi_i \quad (2)$$

which we call the triangular model. Here, $i = 2x/(a\sqrt{3})$ and σ is the dimensionless magnetic flux in units of Bohr magneton, $\sigma = (eH/hc)a^2$ and $\lambda = t_1/t_2$. The isotropic triangular lattice case $t_1 = t_2$ was studied by Claro and Wannier [4] where the model was found to exhibit the hierarchical structure in the energy spectrum similar to that of the square lattice case studied by Hofstadter [5].

In this paper, we show that as a function of λ , the model exhibits a phase transition from critical (C) to localized (L) states. This is in contrast with the case of the square lattice where the anisotropic model exhibits a transition from extended (E) to localized states. Therefore, unlike the square lattice case, the model does not exhibit the E phase of KAM type and hence the weak coupling limit of the model is nontrivial. Furthermore, the critical phase exists in a finite parameter interval. We use our recently developed decimation method [3,6] to study the scaling properties in various parameter ranges. At the band edges, the universality classes of the model are described by the following three non-trivial fixed points of the renormalization operator (the term "fixed point" is used also for periodic cycles of the renormalization operator as the latter are fixed points of an higher-order iteration of the operator): 1) the weak coupling (subcritical) fixed point ($\lambda < 1$), 2) the critical fixed point corresponding to $\lambda = 1$ describing the onset of the C-L transition, and 3) the strong coupling (supercritical) fixed point ($\lambda > 1$) describing the fluctuations in the exponential decay in a localized wave function [7].

The C-L transition and the universal properties of the triangular model (2) are compared with those of the quantum Ising model (QIM) in a transverse field h_i for which the Hamiltonian reads

$$H = - \sum_i [\sigma_i^x \sigma_{i+1}^x + 2h_i \sigma_i^z]. \quad (3)$$

Using the methods described by Lieb et al. [8], the eigenvalue equation for the spin problem can be written in a TBM form, which for $h_i = \lambda \cos[\pi(i\sigma + \phi)]$ becomes

$$\begin{aligned} & \cos[\pi(i\sigma + \phi)]\psi_{i+1} + \cos[\pi(i\sigma + \phi)]\psi_{i-1} \\ & + \lambda \cos[2\pi(i\sigma + \phi)]\psi_i = E\psi_i \end{aligned} \quad (4)$$

Here $E = -\lambda + (\bar{E}^2/4 - 1)/(2\lambda)$, where $\bar{E}$ is the energy of the Ising model (3). [9]

The QIM was recently studied for σ equal to the inverse golden mean $\sigma = (\sqrt{5} - 1)/2$. The model exhibits critical states for $\lambda \leq 1$ where $\lambda = 1$ is the localization transition point. In the QIM, this transition has an additional significance as it also corresponds to the magnetic transition to the long range order. At this point, the lowest energy quantum state is believed to be conformally invariant. [2] Our recent decimation studies [3] confirmed this phase diagram of the model and showed that the subcritical regime was described by a unique fixed point of the renormalization flow. However, at the onset of localization, the renormalization flow was attracted by a different fixed point resulting in a different universality class.

It is rather interesting to note the similarities between these two seemingly unrelated problems. Both are described by a nearest-neighbor TBM where the diagonal as well as the off-diagonal terms are modulating: the periodicity of the diagonal term is twice the periodicity of the off-diagonal term. The only difference between these two models is the fact that unlike the QIM, the triangular model shows a relative phase difference between the diagonal and off-diagonal term which is σ dependent.

We applied the renormalization approach [3] to the triangular model. Our main focuss is to compare the phase diagram and the universal properties of the triangular model with those of the QIM. The TBM is written in the following form, where all sites of the lattice except those labelled by the Fibonacci numbers F_n are decimated:

$$f_n(i)\psi(i + F_{n+1}) = \psi(i + F_n) + e_n(i)\psi(i). \quad (5)$$

The additive property $F_{n+1} = F_n + F_{n-1}$ provides exact recursion relations for the decimation functions e_n and f_n : [3,7]

$$e_{n+1}(i) = -\frac{Ae_n(i)}{1 + Af_n(i)} \quad (6)$$

$$f_{n+1}(i) = \frac{f_{n-1}(i + F_n)f_n(i + F_n)}{1 + Af_n(i)} \quad (7)$$

$$A = e_{n-1}(i + F_n) + f_{n-1}(i + F_n)e_n(i + F_n).$$

In this renormalization approach, the C phase manifests itself in a nontrivial limit cycle of period p , which is typically a multiple of 3, for the absolute values of decimation functions e_n and f_n as $n \rightarrow \infty$. This results in nontrivial scaling ratios

$$\zeta_i = \lim_{n \rightarrow \infty} |\psi(F_{i+pn})/\psi(0)|. \quad (8)$$

Here it is assumed that the phase ϕ is chosen so that the main peak of the wave function lies at $i = 0$. In contrast, the scaling ratios approach unity in the E phase.

In the L phase, we write the wave function as [7],

$$\psi_i = e^{-\gamma|i|}\eta_i \quad (9)$$

where γ is the Lyapunov exponent which vanishes in the E and C phase. η describes the fluctuations in the exponentially decaying part of the wave function. The spatial dependence of η is given by a TBM [7] which is studied by the decimation scheme described above. For the triangular lattice, $\gamma = \ln(\lambda)$ [4]. Due to lack of duality, an analogous expression has not been derived analytically for the QIM. However, a numerical computation of the Lyapunov exponent [10] is found to be in agreement with this formula.

Detailed study of the renormalization flow shows that for the triangular model (2), the decimation functions approach asymptotically the same 3-cycle for both the upper and the lower band edge (maximum and minimum energy). For the QIM, this identity between the scaling properties of the upper and lower band edge is true only in the subcritical and supercritical regimes but not at the critical point (see next paragraph). Furthermore, in both subcritical and supercritical regime, the triangular model and the QIM are described by the same universality class, i.e. the scaling ratios ζ for both models are identical. In the triangular model the wave function has exact symmetry at the band edges whereas in the QIM the wave function is only asymptotically symmetric.

At the critical point $\lambda = 1$ corresponding to the onset of localization, the universal characteristics of the triangular lattice band edges are the same as those of the upper band edge of the QIM. However, the lower band edge of the QIM has zero energy and the corresponding state is believed to be conformally invariant. Unlike the upper band edge 3-cycle, the renormalization flow for the decimation functions at $\bar{E} = 0$ converges to a period-1 fixed point. In addition, the wave function at the conformal point is asymmetrical vanishing on one side of main peak. [3] The "conformal" renormalization fixed point of the QIM does not map to any quantum state of the triangular model.

Fig. 1 shows the self-similar wave functions in the three universality classes for the triangular model at the upper band edge. They also describe the asymptotic wave functions at the upper band edge of the QIM. The dominant peaks in the wave functions are labelled by the rational approximants of σ^3 and its harmonics. [3] The height of these peaks with respect to the central peak define universal scaling ratios. For example, the universal scaling ratios of the peaks at three successive Fibonacci numbers in the subcritical, critical, and supercritical phases are (0.825, 0.908, 0.836), (0.238, 0.303, 0.291), and (0.267, 0.311, 0.121), respectively (see Fig. 1). These scaling ratios are identical to the corresponding scaling ratios at the lower band edge with the exception of the the conformal point in the QIM characterized by the fixed point scaling 0.415.

The study of the band center indicates how the relative phase difference in the off-diagonal terms of the triangular model and the QIM can lead to subtle symmetry properties in the asymptotic renormalization dynamics of the two models. At the band center with $\lambda = 1$, the wave functions are asymmetric in both systems leading to different 6-cycles and scaling ratios (also different from those characterizing the band edges) on the positive and negative side of the lattice. [11] The scaling properties on the positive (negative) side of the triangular model are asymptotically the same as those of the QIM on the negative (positive) side. However, the correspondence is seen only with the shift of three decimation levels between the two systems. Moreover, the critical value of the phase ϕ setting the main peak at the origin is different for the models. In the QIM it is $1/4$ (the same as at the lower band edge) as for the critical model the critical phase is $3/8$ which is the arithmetic mean of the critical phases at the upper and lower band edges.

The Harper equation, which describes Bloch electrons on a square lattice, also models the isotropic XY model in a quasiperiodic field. [3] It is rather interesting that the problem for the Bloch electrons on a triangular lattice is related to the fully anisotropic Ising limit of the XY model. Although the correspondence between the TBM for the square lattice and the isotropic XY model is exact, relationship between the triangular lattice and the Ising model is through the asymptotic universal scaling properties of the models.

Another interesting aspect of our studies is that unlike the periodic case, details of the lattice is relevant in determining the phase diagram and universality classes in incommensurate systems. For the square lattice, there is a phase transition from E to L phase while in the case of a triangular lattice, the C-L transition is observed instead. Therefore, for the square lattice, the weak coupling fixed point is trivial while for the triangular case, the weak coupling is described by the nontrivial fixed point and scaling ratios. Furthermore, even in the L phase, the scaling properties of the self-similar fluctuations in the square and the triangular lattices are different.

The crossover from E-L to C-L phase diagram, as we change from the square to the triangular lattice, can be understood due to the relationship between the 2-dimensional Bloch and spin problem. Previous studies [9] have shown that in the presence of anisotropy, the E and the L phases of the isotropic XY model are mediated by a critical phase with self-similar characteristics. In the Ising limit, the width of the E phase shrinks to zero resulting in the C-L phase diagram. We like to mention that although the Harper equation with next-nearest-neighbor interaction [11,12] provides a means to study the cross-over from square to triangular lattice, understanding the E-L to C-L behavior is lot more complicated.

The results of the triangular model may be of rel-

evance in experiments on quantum dots. [13] Furthermore, since the one-dimensional QIM also describes the two-dimensional classical Ising model with modulating exchange along one direction, it could be realized in experiments on magnetic superlattices. Our studies provide interesting mappings between the thermodynamical properties such as the specific heat and the susceptibility between these two models.

ACKNOWLEDGMENTS

The research of IIS is supported by a grant from National Science Foundation DMR 093296. IIS would like to acknowledge the hospitality of National Institute of Standard and Technology where part of this work is done. JAK has been supported financially by the Academy of Finland under contract 4385 and by the Niilo Helander Foundation. JAK is also grateful for the hospitality of the Research Institute for Theoretical Physics in Helsinki.

-
- [1] For a review, see J. B. Sokoloff, Phys. Rep. 126, 189 (1985).
 - [2] P. Christe and M. Henkel, Introduction to conformal invariance and its applications to critical phenomena. Springer, Berlin 1993.
 - [3] J. A. Ketoja and I. I. Satija, Phys. Lett. A 194, 64 (1994); Renormalization approach to quasiperiodic quantum spin chains, to appear in Physica A.
 - [4] F. H. Claro and G. H. Wannier, Phys. Rev. B 19, 6068 (1979); D. J. Thouless, Phys. Rev. B 28, 4272 (1983).
 - [5] D. R. Hofstadter, Phys. Rev. B 14, 2239 (1976).
 - [6] J. A. Ketoja, Phys. Rev. Lett. 69, 2180 (1992).
 - [7] J. A. Ketoja and I. I. Satija, Self-similarity and localization, (preprint) 1995.
 - [8] E. Lieb, T. Schultz, and D. Mattis, Ann. Phys. (N.Y.) 16, 407 (1961).
 - [9] I. I. Satija, Phys. Rev. B 48, 3511 (1993); Phys. Rev. B 49, 3391 (1994); I. I. Satija and J. C. Chaves, Phys. Rev. B 49, 13239 (1994).
 - [10] We thank J.C. Chaves for confirming this.
 - [11] J. A. Ketoja, I. I. Satija, and J. C. Chaves, Decimation studies of Bloch electrons in a magnetic field: higher order limit cycles underlying the phase diagram, Phys. Rev. B, in press.
 - [12] J. H. Han, D. J. Thouless, H. Hiramoto, and M. Kohmoto, Phys. Rev. B 50, 11365 (1994).
 - [13] R. R. Gerhardts, D. Weiss, and K. Von Klitzing, Phys. Rev. Lett. 62, 1173 (1989).

FIG. 1. The wave function in the subcritical (a), critical (b), and in the supercritical (c) triangular model for the golden mean incommensurability at the upper band edge with $\phi = 0$. Note that in the supercritical case the figure describes the fluctuations $|\eta_i|$ and not $|\psi_i|$.

Fig1 Ketoja and Satija




Fig1 Ketoja and Satija




Fig1 Ketoja and Satija

